

Common fixed-point theorem for commuting maps on a metric space

Suresh Kumar Sahani¹, Vijay Vir Singh², Krishnapal Singh Sisodia³, Kusum Sharma⁴

¹Department of Science and Technology, Rajarshi Janak University, Janakpurdham, Nepal

²Department of Mathematics, School of Basic and Applied Sciences, Lingaya's Vidyapeeth, Faridabad, Haryana, India

³Department of Mathematics, Mody University of Science and Technology, Lakshmangarh, Sikar, Rajasthan, 332311, India

⁴Department of Mathematics National Institute of Technology, Uttarakhand, Srinagar Pauri Garhwal, 246174, India

¹Corresponding author

E-mail: ¹sureshsahani@rju.edu.np, ²singh_vijayvir@yahoo.com, ³sisodiakps@gmail.com, ⁴kusum31sharma@rediffmail.com

Received 15 April 2024; accepted 11 June 2024; published online 20 July 2024
DOI <https://doi.org/10.21595/mme.2024.24160>



Copyright © 2024 Suresh Kumar Sahani, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. Several novel uses of theorems for fixed points in commuting mapping in a fully metric domain are presented. Several conclusions from full metric fixed point theory are improved and extended by our work. Our proofs are inspired by the study of commuting mappings [B. Fisher and S. Sessa, on a fixed point theorem of Gregus, 1986] and [P. Sumati Kumari, Fixed and periodic point theory in certain spaces, 2013].

Keywords: metric space, Cauchy sequence, common fixed-point theorem.

1. Introduction

The concept of a fixed point is essential to a number of different branches of mathematics. many fascinating and interesting branches of mathematics. The search for fixed points is under the purview of many subfields of analysis, such as classical analysis, operator theory, topology, functional analysis, topological algebra, and algebraic topology. Numerous fields make use of fixed-point theorems. different domains, including non-linear oscillations. Theoretical approximation, initial and boundary value difficulties, and the third flow for ordered and partial differential equations. In 1912, Brouwer [1] demonstrated that the mapping is continuous ζ in reference to the closed unit ball R^1 space having at least one fixed point, which we'll refer to as point z : a point where $\zeta_z = z$. In 1922, mathematicians Birkhoff and Kellogg [2] developed the first theory of fixed points in infinite dimensions. Using Brouwer's Birkhoff and Kellogg provided evidence for the existence of theorems in the theory of differential equations through their work on the fixed-point theorem. In the case where E is a convex compact set that exists in a normed vector space, Schauder [3, 4] generalized Brouwer's fixed point theorem in 1927 and 1930, respectively. Tychonoff [5] was the first mathematician to generalize the Schauder result from normed space to locally convex topological vector space considered by the authors in [35]. It was first proven in 1986 by Fisher and Sessa [6] that two self-maps have a fixed point if they are located on a convex subset of a Banach space. Refer to the work Sessa, [7] elaborated in which Das&Naik, [8] research was represented as a reference. Jungeek [9] first generalized the Sessa concept to compatible mapping, and later to weakly compatible mapping (see [10]). Several authors have extended Jungeek's approach to produce coincidence point findings for various types of mapping on Fatus-proper metric spaces (see [11,12]). Some generalizations points of coincidental occurrence and popular fixed points in fixed point theory were shown to be false by Haghi et al. (2011) in which they could be produced immediately from the fixed-point theorems that are linked to it.

Jleli and Samet [13] were the ones who came up with the idea of a generalized metric space,

which has subsequently been used to the process of recovering a number of different topological spaces. Metric, b-metric, displaced, and modular spaces are examples (see [16-24, 30]).

Using the E. A property on metric space, authors L. Wangwe and Santosh Kumar [25] studied the common fixed-point theorem under implicit contractive conditions. An arbitrary binary relation has been developed and applied to second-order differential equations with boundary conditions. In his study of fixed point and continuity for a pair of contractive maps, Santosh Kumar [26] applied the nonlinear Volterra Integral equation and provided multiple examples proving the existence and uniqueness of the solution. In TVS valued cone metric space, Lucus Wangwe and Santosh Kumar [27] have investigated to generalize and extend several fixed points results that illustrate the common fixed points theorem for F-Kanan- Suzuki type mapping. Dur-e-Shehwar Sangheer et al. [28] have investigated a novel multi valued F- contraction mapping incorporating α - admissibility with applications, where they have proved on existence solution of integral equations. Recently authors Lucus Wangwe [29] proved the common fixed points theorem for interpolative rational type of contraction mapping in complex valued metric space with application in existence and uniqueness solution of R-L-C differential equation.

Definition [14]: Take the set F to be non-empty. Where $m, n \in F$ and (F, d) be metric space. A mapping $\xi: F \rightarrow F$ is said to be a generalized contraction if we find a positive real number δ_1 satisfies the following conditions:

$$d(\xi m, \xi n) \leq \delta_1 \max \{d(m, n), d(m, \xi m), d(n, \xi n), d(m, \xi n) + d(n, \xi m)\}. \quad (1)$$

Known theorems: In 1979, Fisher [15] the following theorem.

Theorem: Let (R, d) be entire metric space, ξ and ζ are continuous functions (signals) of (R, d) into itself. Then both signals have a common stationary in the set R if only if we find a continuous signal F of R into $\xi(R) \cap \zeta(R)$ commutes with two signals G and ζ meet certain requirements $F(R) \subset \xi(R) \cap \zeta(R)$, $d(Fm, Fn) \leq \delta d(\xi m, \zeta n)$, where, $m, n \in R$, $\delta \in (0,1)$, in fact, G and ζ having a single point of agreement.

Let's say that ξ , ζ , and F are three self-mapping of R satisfying the following constraints:

$$F\xi = \xi F, \quad (2)$$

$$F\zeta = \zeta F, \quad (3)$$

$$F(R) \subset \xi(F) \cap \zeta(F), \quad (4)$$

where ξ and ζ are continuous. For all $m, n \in F$ we find a positive integer δ_1 such that:

$$d(Fm, Fn) \leq \delta_1 \max \left\{ d(\xi m, \zeta n), d(\xi m, Fm), d(\zeta n, Fn), \frac{d(\xi m, Fn), d(\zeta n, Fm)}{2} \right\}. \quad (5)$$

Lemma: If F , ξ , and ζ are self-mapping of R satisfying when (2), (3), and (4) hold, then is what R calls a Cauchy sequence.

Proof: Let m_o be a random number in R . Given that $F(R)$ is encapsulated in $\xi(R)$, we choose a special point m_1 in R such that $\xi m_1 = Fm_o$.

Since $F(R)$ is also contained in $\zeta(R)$, we choose another special point m_2 in F such that $\xi m_2 = Fm_1$. In the same way, we choose a point m_r such that $\xi m_r = Fm_{r-1}$ when r is even.

Now we have to show that $\{Fm_r\}$ is a Cauchy sequence in R . Using the Eq. (5) we obtain:

$$\begin{aligned} d(Fm_1, Fm_2) &\leq \delta_1 \max \left\{ d(\xi m_1, \zeta m_2), d(\xi m_1, Fm_2), d(\zeta m_2, Fm_2), \frac{d(\xi m_1, Fm_2), d(\zeta m_2, Fm_2)}{2} \right\} \\ &= \delta_1 \max \left\{ d(Fm_o, Fm_1), d(Fm_1, Fm_2), \frac{1}{2}d(Fm_o, Fm_2) \right\}. \end{aligned} \quad (6)$$

Thus, we obtain $d(Fm_1, Fm_2) \leq d(Fm_o, Fm_1)$.

In the same way, we can easily obtain $d(Fm_r, Fm_{r+1}) \leq \delta_1^r d(Fm_o, Fm_1) \Rightarrow \{Fm_r\}$ for all $r \in N$, contains a Cauchy sequence in R .

Theorem 1: Allow a whole metric domain to map itself using F , ξ , and ζ If (2), (3), (4), and (6) hold, if this is the case, then F , ξ , and ζ all have a common fixed point.

Proof: Suppose m_o is a chosen coordinate in R , and $\{Fm_r\}$ be a sequence satisfying all conditions of the given lemma. Then $\{Fm_r\}$ transforms into a Cauchy sequence. There is just one possible limit point for a convergent Cauchy sequence. Let us suppose that $\{Fm_r\} \rightarrow l$. Since the sequences, $\{\xi m_{2r+1}\}$ and $\{\zeta m_{2r}\}$ are the subsequences of $\{Fm_r\}$. We know that every subsequence of a convergent sequence has the same limit point. Thus, we have $\{G\xi\} \rightarrow l$ and $\{\zeta m_{2r}\} \rightarrow l$ as $r \rightarrow \infty$.

Consequently, we write $\zeta(Fm_r) \rightarrow \zeta l$, $\xi Fm_{2r+1} \rightarrow \xi l$, $\zeta(\xi m_{2r+1}) \rightarrow \zeta l$, and $\xi(\zeta m_{2r}) \rightarrow \xi l$. We consider:

$$d(\zeta Fm_{2r}, \xi Fm_{2r+1}) = d(F\zeta m_{2r}, F\xi m_{2r+1}) \\ \leq \delta_1 \max \left\{ d(\xi \zeta m_{2r}, \zeta \xi m_{2r+1}), d(\xi \zeta m_{2r}, F\zeta m_{2r}), d(\zeta \xi m_{2r+1}, F\xi m_{2r+1}), \right. \\ \left. d(\zeta \xi m_{2r}, F\xi m_{2r+1}), \frac{1}{2} (d(\zeta \xi m_{2r}, F\xi m_{2r+1}), d(F\zeta m_{2r}, \zeta \xi m_{2r+1})) \right\}$$

Using $r \rightarrow \infty$ then $d(\zeta l, \xi l) \leq \delta_1 \max\{d(\xi l, \zeta l), d(\xi l, \zeta l), d(\zeta l, \xi l)\} \Rightarrow \zeta l = \xi l$.
Again, from the Eq. (5):

$$d(\zeta Fm_{2r}, Fl) = \left\{ d(F\zeta m_{2r}, Fl), d(\xi \zeta m_{2r}, \zeta l), d(\zeta l, Gl), \frac{1}{2} (d(\xi \zeta m_{2r}, Fl), d(F\zeta m_{2r}, Fl)) \right\}$$

Using $\rightarrow \infty$, we get $d(\zeta l, Fl) \leq d(\zeta l, Fl) \Rightarrow \zeta l = Fl$.
Thus, $Fl = \zeta l = Gl$.
Next, we consider:

$$d(\zeta m_{2r}, Fl) = d(Fm_{2r-1}, Fl) \leq \max \left\{ d(\xi m_{2r-1}, \zeta l), d(\xi m_{2r-1}, Fm_{2r-1}), \right. \\ \left. d(\zeta l, Fl), \frac{1}{2} (d(\xi m_{2r-1}, Fl), d(Fm_{2r-1}, \zeta l)) \right\}$$

Using $r \rightarrow \infty$, we obtain $d(l, Fl) \leq \delta_1 d(l, Fl) \Rightarrow l = Fl$.

At every condition, we have seen that l is a fixed point of ξ , F , and ζ .

Theorem 2: Assume the following about the compact metric space (R, d) : F , ξ , and ζ are all self-mappings satisfying:

$$d(Fm, Fn) < \max \left\{ d(\xi m, Fn), d(\xi n, \zeta n), d(\zeta n, Fm), \frac{d(\xi m, Fn) + d(\zeta n, Fm)}{2} \right\}, \quad (7)$$

for all $m, n \in R$ and R.H. S of Eq. (7) is a positive number.

If the self-mapping F , ξ and ζ are continuous then they have a unique common fixed point.

Proof: Let us Let's pretend the answer to the right side of the problem is Eq. (7) is positive. Then we define a function $\eta(m, n)$ be defined by:

$$\eta(m, n) = \frac{d(Fm, Fn)}{\max\{d(\xi m, Fm), d(\xi m, \zeta m), d(\zeta n, Fn), \frac{1}{2}(d(\xi m, Fn) + d(\zeta n, Fm))\}}$$

is continuous real valued mapping on the compact metric space and attains a maximum value of δ . From the Eq. (7) we choose a number δ such that $\delta > 0$ and hence by using theorem 1, the mapping F , ξ and ζ have a single, unifying point of reference.

Let's pretend the answer to Eq. (7) on the right is zero since $m, n \in R$.

Thus $\xi m = \zeta n = Fm = Fn \Rightarrow F\xi m = F^2m = \xi Fm$.

If $F^2m \neq Fn$ then:

$$d(F^2m, Fn) < \max \left\{ d(\xi Fm, F^2m), d(\xi Fm, \zeta n), d(\zeta n, Fn), \frac{1}{2}[d(\xi Fm, Fn) + d(\zeta n, F^2m)] \right\}$$

$$\therefore d(F^2m, Fn) < d(F^2m, Fn),$$

which gives a contradiction.

Thus, $F^2m = Fn = F(Fm) = F(Fn)$. $\therefore Fn = l$ is a constant value that F .

And $\xi l = \xi(Fn) = \xi(Fm) = F(\xi m) = F(Fn) = Fl = l$, and $\zeta l = \zeta(Fn) = F(\zeta n) = F(Fn) = Fl = l$.

As a result, we can declare that l is the place where F , ξ , and ζ all meet.

Corollary 1: Let F and ζ be compact metric (R, d) mappings that commute into themselves such that:

$$d(\zeta n, Fn) < \max \left\{ d(\zeta m, \zeta n), d(\zeta m, Fm), d(\zeta n, Fn), \frac{d(\zeta m, Fn) + d(\zeta n, Fm)}{2} \right\}$$

holds for all $m, n \in R$ for which its R.H.S. is non-negative.

If $F(R) \subset \zeta(R)$, F and ζ is an extended result of the authors [15] and states that if two maps are continuous, then they must share a fixed point.

Corollary 2: Let F be a continuous function of a compact metric space (R, d) into itself such that:

$$d(Fm, Fn) < \max \left\{ d(m, n), d(m, Fm), d(n, Fn), \frac{d(m, Fn) + d(n, Fm)}{2} \right\}$$

holds for all $m, n \in R$ for which its R. H.S. of above inequality is non-negative, if F has a shared fixed point, then. (This finding generalizes the author's previous findings; see [16]).

Corollary 3: If ζ a continuous function of a compact metric space (R, d) into itself which satisfy the following condition:

$$d(m, n) < \max \left\{ d(\zeta m, \zeta n), d(\zeta m, m), d(\zeta n, n), \frac{d(\zeta m, n) + d(\zeta n, m)}{2} \right\},$$

$\forall m, n \in R$ for which its R. H. S of its inequality is non-negative then the continuous mapping ζ has a unique fixed point.

Proof: If F be a continuous identity mapping and $F(R) \subset \zeta(R)$. With the help of corollary Eq. (2), we get our result.

Theorem 3: Let F , ξ , and ζ be self-mappings of a complete metric space which satisfy the following conditions:

- 1) $F\zeta = \zeta F, F\xi = \xi F$.
- 2) $F(R) \subset \zeta^a(R) \cap \xi^b(R)$.
- 3) ξ^b and ζ^a are continuous.
- 4) F^c, ξ^b and ζ^a satisfy:

$$d(F^c m, F^c n) < \delta_1 \max \left\{ d(\xi^b m, \zeta^a n), d(\xi^b m, F^c m), d(\zeta^a n, F^c n), \frac{d(\xi^b m, F^c n) + d(\zeta^a n, F^c m)}{2} \right\},$$

where, a and b are some positive integers and $c > 0$ Therefore, F , ξ , and ζ have a common fixed point.

Proof: Since F commutes with the mapping ξ and ζ , F^c commutes with ξ^b and ζ^a . Further $F^c(R) \subset F(R) \subset \xi^b(R)$. In the same line, we may write $F^c(R) \subset \zeta^a(R)$.

Also, since F^c , ξ^b and ζ^a are continuous mapping then by using theorem 1, we find a point $l \in R$ such that $l = F^c l = \xi^b l = \zeta^a l$.

Also, $\xi^a(Fl) = F(\xi^a l) = Fl = F(F^c l) = F^c(Fl) \Rightarrow Fl$ is a common fixed of ξ^b and F^c .

In the same line previous work, we define Fl as the fixed point in common between ζ^a and F^c .

By the definition of uniqueness of a point $l \Rightarrow l = Fl$.

Let $\xi^{b+1}l = \xi l$ and $\zeta^a l = \zeta l$ and if we substitute $m = Fl$ and $n = Fn$ in Eq. (5) we easily obtain $\xi l = \zeta l \therefore \xi l = \zeta l$ is also a common point for F^c , ξ^b and ζ^a .

Thus, by uniqueness of common fixed point l , we obtain $l = Fl = \xi l = \zeta l$.

This result extends the result of author [15].

Some Applications. If we slight change in our theorems then we can easily obtain same famous results of periodic point theory in certain spaces. For this, we study.

Definition 1: Assume ξ is a self-map of R , and $m \in R$ and $a \in N$. If $\xi^a m = m$ then m is known as periodic point of the function ξ .

Definition 2: Let $m \in R$, $a, b \in N$, and ξ and ζ are both self mappings of a complete metric space (R, d) . If $\xi^a m = \zeta^a m = m$ then a given point m is known as common periodic point of our function ξ and ζ .

Definition 3: Given a non-empty set R , we say that $m, n, l \in R$. A metric for a collection that is not empty as a function, $Rd: R \times R \rightarrow [0, \infty)$ such that:

- 1) $d(l, l) = 0$.
- 2) $d(l, m) = d(m, l) = 0 \Leftrightarrow m = l$.
- 3) $d(l, m) = d(m, l)$.
- 4) $d(l, m) \leq d(l, n) + d(n, m)$.

Theorem 4: Let ξ and ζ are continuous functions of complete metric space (R, d) satisfying:

$$d(\xi^b, \zeta^a n) < \max \left\{ d(m, n), \frac{d(\xi^b m, m) + d(\zeta^a n, n)}{2}, \frac{d(\zeta^a n, m) + d(\xi^b m, n)}{2} \right\}, \quad (8)$$

for all $m, n \in R$ ($m \neq n$) in which R.H. S. of Eq. (8) is non-zero and $a, b \in N$. Then $r \in R$ is common periodic of mappings ξ and ζ point if and only if r is the single fixed point in both mappings.

Proof: It is obvious that only any periodic point is referred to as a fixed point. For conversely, we put $\xi^b = L$ and $\zeta^a = M$ in our Eq. (8) then we have:

$$d(Lm, Mn) < \max \left\{ \frac{d(m, n) + d(Lm, m) + d(n, Mn)}{2}, \frac{d(Mn, l) + d(n, Lm)}{2} \right\}.$$

Setting $a = b = 1$ in the theorem Eq. (8) we represent the following is an illustration.

Let $R = [0, 1)$ and $d(m, n) = |m - n|$. Let $G, \zeta: R \rightarrow R$ by $Gm = m/3$ and $\zeta m = 0$.

The value 0 is the only common fixed point that exists. $\Rightarrow 0$ is the unique common periodic point.

Here ξ and ζ satisfy the aforementioned inequality, and verification may be obtained by looking at the following:

- 1) $m = 0, 0 < n < 1 \therefore$ L.H.S. = 0 and R. H. S. = $|n|$.
- 2) $n = 0, 0 < m < 1$, L.H.S = $|m|/3$ and R. H. S. = $|m|$.
- 3) for $1 > n > m > 0$ (or for $1 > m > n > 0$) we obtain L.H.S and R.H.S. = $\max\{|m -$

$n|, |2m| + |n|/3, |m| + |n + \frac{m}{3}|/2\}$.

Thus L.H.S. = $|m|/3 < \frac{|2m|/3 + |n|}{2} \leq$ R.H.S.

The self-mappings ξ and ζ satisfy above inequality when $a = 2$ and $b = 3$.

Theorem 5: Let ξ and ζ are continuous function of complete metric space (R, d) satisfying:

$$d(\xi^b m, \zeta^a n) < \max \left\{ d(m, n), \frac{d(\xi^b m, m) + d(\zeta^a n, n)}{2}, \frac{d(\zeta^a n, m) + d(\xi^b m, n)}{2} \right\}, \quad (9)$$

for all $m, n \in R$ ($m \neq n$) in which R.H.S. of Eq. (9) is non- zero and $a, b \in N$. Then $r \in R$ is r is common periodic point of mappings if and only if r is considered to be the common fixed point of mappings ξ and ζ .

Proof: If we put $b = a$ in the given theorem Eq. (8), we easily obtain above theorem 5.

Theorem 6: Consider the function ξ which is continuous mapping from a complete metric space (R, d) into itself satisfying:

$$d(\xi^b m, \xi^a n) < \max \left\{ d(m, n), \frac{d(\xi^b m, m) + d(\xi^a n, n)}{2}, \frac{d(\xi^b m, n) + d(\xi^a n, m)}{2} \right\}, \quad (10)$$

for all $m, n \in R$ ($m \neq n$) and $a, b \in N$ in which R.H.S of Eq. (10) is non-negative numbers. Then $r \in R$ is unique fixed point of the function the function ξ if u is a periodic point of the function ξ .

Proof: If we put $\zeta = \xi$ in Eq. (8), we can readily establish theorem Eq. (10).

Theorem 7: Let ξ be a continuous function a complete metric space (R, d) into itself satisfying:

$$d(\xi^b m, \xi^b n) < \max \left\{ d(m, n), \frac{d(\xi^b m, m) + d(\xi^b n, n)}{2}, \frac{d(\xi^b n, m) + d(\xi^b m, n)}{2} \right\}, \quad (11)$$

for all $m, n \in R$ ($m \neq n$) and $b \in N$ in which R.H S of Eq. (11) is non-negative numbers. Then $r \in R$ is unique fixed point of the function ξ if and only if r is the periodic point of function ξ .

Proof: Put $a = b$ in theorem Eq. (10), we get the proof of the theorem 7.

Acknowledgements

The authors have not disclosed any funding.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Author contributions

Dr. Suresh Kumar Sahani contributed as author for preparing the manuscript and edited the article incorporated the comments of reviewers. Dr. Sahani, (Prof). Vijay Vir Singh, Dr. Krishnapal Singh Sisodia, and Dr. Kusum Sharma have revised the manuscript with their own insights to submitting the final revision.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] L. E. J. Brouwer, "Über abbildung von mannigfaltig keiten," *Mathematische Annalen*, Vol. 71, No. 4, p. 598, Dec. 1912, <https://doi.org/10.1007/bf01456812>
- [2] G. D. Birkhoff and O. D. Kellogg, "Invariant points in function space," *Transactions of the American Mathematical Society*, Vol. 23, No. 1, pp. 96–115, Jan. 1922, <https://doi.org/10.1090/s0002-9947-1922-1501192-9>
- [3] J. Schauder, "Zur Theorie stetiger Abbildungen," *Jahresbericht der Deutschen Mathematiker-Vereinigung*, Vol. 59, pp. 131–140, 1957.
- [4] J. Schauder, "Zur Theorie stetiger Abbildungen, der Fixpunktsatz in Funktionalräumen," *Studia Mathematica*, Vol. 2, pp. 171–180, 1930.
- [5] A. Tychonoff, "Fest Fixpunkte," *Mathematische Annalen*, Vol. 111, pp. 767–776, 1935.
- [6] B. Fisher and S. Sessa, "On a fixed point theorem of Greguš," *International Journal of Mathematics and Mathematical Sciences*, Vol. 9, No. 1, pp. 23–28, Jan. 1986, <https://doi.org/10.1155/s0161171286000030>
- [7] S. Sessa, "On a weak commutativity condition of mappings in fixed point considerations," *Publications de l'Institut Mathématique*, Vol. 32, No. 52, pp. 149–153, 1982.
- [8] K. M. Das and K. V. Naik, "Common fixed-point theorems for commuting maps on a metric space," *Proceedings of the American Mathematical Society*, Vol. 77, No. 3, pp. 369–373, Jan. 1979, <https://doi.org/10.1090/s0002-9939-1979-0545598-7>
- [9] G. Jungck, "Compatible mappings and common fixed points," *International Journal of Mathematics and Mathematical Sciences*, Vol. 9, No. 4, pp. 771–779, Jan. 1986, <https://doi.org/10.1155/s0161171286000935>
- [10] G. Jungck, "Common fixed points for non-continuous nonself maps on nonmetric spaces," *Far East Journal of Mathematical Sciences*, Vol. 4, No. 2, pp. 199–215, 1996.
- [11] G. Jungck and N. Hussain, "Compatible maps and invariant approximations," *Journal of Mathematical Analysis and Applications*, Vol. 325, No. 2, pp. 1003–1012, Jan. 2007, <https://doi.org/10.1016/j.jmaa.2006.02.058>
- [12] G. Jungck, "Common fixed points for commuting and compatible maps on compacta," *Proceedings of the American Mathematical Society*, Vol. 103, No. 3, pp. 977–983, Jan. 1988, <https://doi.org/10.1090/s0002-9939-1988-0947693-2>
- [13] M. Jleli and B. Samet, "A generalized metric space and related fixed point theorems," *Fixed Point Theory and Applications*, Vol. 2015, No. 1, Apr. 2015, <https://doi.org/10.1186/s13663-015-0312-7>
- [14] Lj. B. Ćirić, "Generalized contractions and fixed point theorems," *Publications de l'Institut Mathématique*, Vol. 26, pp. 19–26, 1971.
- [15] G. Jungck, "An iff fixed point criterion," *Mathematics Magazine*, Vol. 49, No. 1, pp. 32–34, Feb. 2018, <https://doi.org/10.2307/2689880>
- [16] M. Edelstein, "On fixed and Periodic points under contractive mappings," *Journal of the London Mathematical Society*, Vol. s1-37, No. 1, pp. 74–79, Dec. 2016, <https://doi.org/10.1112/jlms/s1-37.1.74>
- [17] K. Chaira, M. Kabil, and A. Kamouss, "Common Fixed Points in Generalized Metric Spaces with a Graph," *Journal of Mathematics*, Vol. 2019, pp. 1–8, Feb. 2019, <https://doi.org/10.1155/2019/2846315>
- [18] J. J. Nieto and R. Rodríguez-López, "Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations," *Order*, Vol. 22, No. 3, pp. 223–239, Nov. 2005, <https://doi.org/10.1007/s11083-005-9018-5>
- [19] K. Chaira et al., "Some generalised fixed point theorems in a partially ordered space endowed with two metrics," *Advances in Fixed Point Theory*, Vol. 6, No. 4, 2017.
- [20] M. R. Alfuraidan, M. Bachar, and M. A. Khamsi, "Almost monotone contractions on weighted graphs," *Journal of Nonlinear Sciences and Applications*, Vol. 9, No. 8, pp. 5189–5195, Aug. 2016, <https://doi.org/10.22436/jnsa.009.08.04>
- [21] W. D. Wallis, *A Beginner's Guide to Graph Theory*. Boston, MA: Birkhäuser Boston, 2000, <https://doi.org/10.1007/978-1-4757-3134-7>
- [22] Z. Mustafa et al., "Some common fixed point results of graphs on b-metric spaces," *Journal of Nonlinear Sciences and Applications*, Vol. 9, No. 6, pp. 4838–4851, 2016.
- [23] P. Sumati Kumari, "Fixed and periodic point theory in certain spaces," *Journal of the Egyptian Mathematical Society*, Vol. 21, No. 3, pp. 276–280, Oct. 2013, <https://doi.org/10.1016/j.joems.2013.04.007>

- [24] M. Abbas and B. E. Rhoades, “fixed and periodic point results in cone metric spaces,” *Applied Mathematics Letters*, Vol. 22, No. 4, pp. 511–515, Apr. 2009, <https://doi.org/10.1016/j.aml.2008.07.001>
- [25] L. Wangwe and S. Kumar, “Common fixed point theorems under implicit contractive condition using E. A. property on metric-like spaces employing an arbitrary binary relation with some application,” *International Journal of Nonlinear Analysis and Applications*, Vol. 13, Jun. 2022, <https://doi.org/10.22075/ijnaa.2021.23494.2548>
- [26] S. Kumar, “Fixed points and continuity for a pair of contractive maps with application to nonlinear volterra integral equations,” *Journal of Function Spaces*, Vol. 2021, pp. 1–13, Aug. 2021, <https://doi.org/10.1155/2021/9982217>
- [27] L. Wangwe and S. Kumar, “Common fixed point theorems for F-Kannan-Suzuki type mappings in TVS-valued cone metric space with some applications,” *Journal of Mathematics*, Vol. 2022, pp. 1–17, Apr. 2022, <https://doi.org/10.1155/2022/6504663>
- [28] D.-E.-S. Sagheer, S. Batul, I. Urooj, H. Aydi, and S. Kumar, “New multivalued F-contraction mappings involving α -admissibility with an application,” *Journal of Inequalities and Applications*, Vol. 2023, No. 1, Aug. 2023, <https://doi.org/10.1186/s13660-023-03016-x>
- [29] L. Wangwe, “Common fixed point theorems for interpolative rational-type mapping in complex-valued metric space,” *European Journal of Mathematics and Applications*, Vol. 4, p. 15, Jan. 2024, <https://doi.org/10.28919/ejma.2024.4.15>
- [30] S. K. Sahani, B. K. Pandey, and D. Pandey, “Single-valued signals, multi-valued signals and fixed-point of contractive signals,” *Mathematics Open*, Mar. 2024, <https://doi.org/10.1142/s2811007224500020>



Dr. **Suresh Kumar Sahani** is an Assistant Professor in the Department of Science and Technology at Rajarshi Janak University, Janakpur, Nepal. He has received his Ph.D. Degree in 2017 in Applied Mathematics from Shri Venkateshwara University, UP, India. His broad research area includes mathematical analysis, summability, approximation theory, and applied mathematics. He has published more than 150 research papers in National and International reputed Journals.



Dr. (Prof.) **Vijay Vir Singh** received his MSc, degree (1989), MPhil degree (1992) & Ph.D. (2010) from Chaudhary Charan Singh University campus Meerut, India. Dr. Singh has worked as Professor in various positions in different countries. Prof. Singh’s research field is in Operation research specialized in reliability theory with copula linguistic approach. He has supervised four Ph.D., Six MSc, and more than 40 students for BSc in reliability and other applied mathematics area, complex analysis (Weierstrass p functions). Dr. Singh has participated in many international conferences in India, Thailand, and Nigeria as a presenter, technical advisory board member, and session chairperson. Prof. V. V. Singh is a member of the editorial boards several Journals in India and abroad. He has reviewed many research articles from Springer, Elsevier, Inderscience, and other SCI journals in Reliability and Applied Mathematics areas. Dr. Singh has published approx. 85 research papers in his specialized area reliability and statistics which are available at Research Gate.



Dr. **Krishnapal Singh Sisodia** is working as an Assistant Professor, Department of Mathematics, School of Liberal Arts and Sciences, Mody University of Science and Technology, Lakshmanagarh, Sikar (Raj.). He completed his Ph.D. from Barkatullah University, Bhopal (M.P.) in the year 2015. His research interests include Functional Analysis, Topology, Integral transform and Algebra. He has published 10 research papers in national and international referred journals, presented 7 papers in national and international conferences and attended several conferences, workshops, seminar, webinar and Faculty Development Programs. He has about 15 years of experience in teaching and research. Google Scholar citations are 9 and h-index is 2.



Dr. **Kusum Sharma** is an Assistant Professor in the Department of Mathematics at the National Institute of Technology, Uttarakhand. She has received her Ph.D. Degree in 2013 in pure mathematics from Mody University of Science and Technology, Rajasthan. Her broad research area includes analysis, operator theory, summability, and approximation theory. She has published more than 18 research papers in journals of international repute